

Emergency treatment of chemical and thermal eye burns

Ralf Kuckelkorn, Norbert Schrage, Gabriela Keller and Claudia Redbrake

Department of Ophthalmology, Universitätsklinikum der RWTH Aachen, Aachen, Germany

ABSTRACT.

Chemical and thermal eye burns account for a small but significant fraction of ocular trauma. The speed at which initial irrigation of the eye begins, has the greatest influence on the prognosis and outcome of eye burns. Water is commonly recommended as an irrigation fluid. However, water is hypotonic to the corneal stroma. The osmolarity gradient causes an increased water influx into the cornea and the invasion of the corrosive substance into deeper corneal structures. We therefore recommend higher osmolarities for the initial rinsing to mobilize water and the dissolved corrosives out of the burnt tissue. Universal systems such as amphoteric solutions, which have an unspecific binding with bases and acids, provide a convenient solution for emergency neutralisation. Both conservative anti-inflammatory therapy and early surgical intervention are important to reduce the inflammatory response of the burnt tissue. In most severe eye burns, tenoplasty re-establishes the conjunctival surface and limbal vascularity and prevents anterior segment necrosis.

Key words: cornea - emergency treatment - eye burns - irrigation fluid - reconstructive surgery.

Acta Ophthalmol. Scand. 2002; 80: 4-10
Copyright © Acta Ophthalmol Scand 2002. ISSN 1395-3907

Recent studies put the incidence of chemical and thermal injuries to the eye at 7.7%–18% of all ocular trauma (Watz & Reim 1973; Pfister et al. 1984; Liggett 1989; MacEwen 1989; Zigelbaum et al. 1993). Most of these injuries are trivial and do not cause any lasting lesions, others result in permanent unilateral or bilateral visual impairment and a life of dependency (Kuckelkorn et al. 1993). The majority of victims are young and exposure occurs at home, work and in association with criminal assaults (Keeney 1974; Morris et al. 1987; Thielsch et al. 1989). Alkali injuries occur more frequently than acid injuries (Pfister 1983; Morgan 1987), eye burns caused by detergents and thermal agents being less frequent again (Kuckelkorn et al. 1995). The most common agents causing alkali burns are ammonia (NH₃), lye (NaOH), potassium hydroxide (KOH) and lime

(CaOH₂). Sulfuric (H₂SO₄), sulfurous (H₂SO₃), hydrofluoric (HF) and hydrochloric (HCl) acids are the most common causes of acid burns. Table 1 lists the data from 191 patients with 260 severely burnt eyes who were treated in the eye center of the RWTH Aachen between 1980 and 1995. There was a high incidence of bilateral injuries and most of the injuries occurred at home or during leisure activities.

Action of alkalis and acids

The severity of ocular injury is related to the type of chemical, the volume and concentration (pH) of the solution and the duration of exposure (Hughes 1946). Alkalis penetrate more rapidly than acids. The hydroxylion (OH⁻) saponifies the fatty acid components of the cell membranes with consecutive cell disruption and cell death, while the cation is respon-

sible for the penetration process of the specific alkali (McCulley 1987). The penetration rate increases from calcium hydroxide (slowest), potassium hydroxide (faster), sodium hydroxide (even faster) to ammonium hydroxide (fastest; Grant 1974). Depending on the degree of penetration, there is a loss of corneal and conjunctival epithelium, stromal keratocytes and endothelium. Hydration of the glycosaminoglycans results in loss of clarity of the stroma (Grant & Kern 1955). Damage to the vascular endothelium of conjunctival and episcleral vessels leads to thrombosis of the episcleral vessels.

The stronger the alkali, the faster its penetration. Irreversible damage occurs at a pH above 11.5 (Friedenwald et al. 1944). The pH in the aqueous humour rises within a few seconds of contact with ammonium hydroxide (Graupner & Hausmann 1970). Intraocular structures such as the iris, lens and ciliary body are rapidly damaged.

Table 1. Severe chemical and thermal eye burns in the Department of Ophthalmology of the RWTH Aachen (1985-1995): 191 patients (260 eyes).

	Number of patients	Per cent
Unilateral	122	64
Bilateral	69	36
Eyes		
Occupational injuries	177	68.1
Private injuries	63	24.1
Others	20	7.8
Eyes		
Alkalis	151	58.1
Acids	37	14.1
Thermal	42	16.2
Others	30	11.6

Apart from hydrofluoric acid and, to a lesser extent, sulfurous acid, acids penetrate the corneal stroma much less readily than alkalis (Grant 1974). The hydrogen ion causes damage due to pH alteration, while the anion produces protein precipitation and denaturation in the corneal epithelium and superficial stroma (Friedenwald et al. 1946). Precipitation of the epithelial proteins offers some protection to the corneal stroma and intraocular structures. However, very strong acids penetrate just quickly as alkalis. No statistical difference between strong alkali and acids burns was discovered in the clinical course and prognosis of such eyes (Kuckelkorn 1996).

Clinical classification of chemical and thermal burns

Eye burns are classified in 4 grades (Reim 1987, 1990). The clinical course and ultimate prognosis correlates with the extent of limbal ischemia (Hughes 1946; Ballen 1963; Roper-Hall 1965). The prognosis also depends on the extent of damage to conjunctival and episcleral tissue, severity of lid burn and damage to intraocular structures (Table 2).

Mild burns of grades I and II are associated with hyperemia, small conjunctival ecchymosis and chemosis as well as erosion of the corneal epithelium (Figs 1 and 2). In mild acid burns, the coagulated corneal epithelium often has a 'ground-glass' appearance. After removal of the epithelium, the clear corneal stroma is visible.

Grade III, and especially grade IV, burns are accompanied by extensive and deep damage to the tissue (Figs 3 and 4). Typically, large areas of the conjunctival and subconjunctival tissue are involved. The visible blood vessels are thrombosed and appear dark. The corneal keratocyt-




Fig. 1. Grade I chemical injury: hydrochloric acid (HCl). Burn of the cornea only. Coagulated corneal epithelium with 'ground glass' appearance. Partial removal of the epithelium, clear corneal stroma.

es are lost and hydration of the denatured proteins results in corneal opacification. Chemical injury to the iris and crystalline lens may produce mydriasis, a greyish appearance of the iris and the fast development of a cataract. The lysis of cells of the anterior chamber destroys the blood aqueous barrier and leads to iridocyclitis and fibrinous exudation.

Toxic substances such as prostaglandins, superoxide radicals, and presumably histamine, angiotensin, leukotriens and others are released from the burnt cells of the necrotic tissue (Eakins & Bhattacharjee 1977; Kulkarni & Srinivasan 1993; Rochels et al. 1982). An inflammatory response is initiated, when they diffuse into surviving tissues. In mild burns this reac-

tion resolves quickly, while in severe burns a severe and long-term inflammatory process is initiated, determining the clinical course of the burnt eyes (Reim 1982, 1987, 1992; Williams et al. 1983; Struck et al. 1991; Reim & Leber 1993; Reim et al. 1993).

Emergency treatment

Immediate irrigation is of paramount importance after chemical or thermal burns (Lubeck & Greene 1988; Cohen & Hyndiuk 1978; Rodeheaver et al. 1982). In most cases the victims are disabled by severe reflexory blepharospasm with ensuing disorientation. In this situation the victims are unlikely to be capable of reaching the nearest body or eye shower and need res-

Table 2. Clinical classification and prognosis of eye burns

Grade	I	II	III	IV
Appearance	Erosio	Erosio	Erosio	Erosio
Limbal ischemia > 3/4	Hyperemia	Limbal ischemia > 1/3 Chemosis	Limbal ischemia > 1/2 Chemosis	Chemosis
Opacification	Opacification	Recirculation	Vascularisation	Extensive necrosis
Clinical outcome	Regeneration	Regeneration	Ulceration Proliferaton Cicatrizaton	Ulceration Iris atrophy Cataract Glaucoma
Prognosis	Complete restitution	Complete restitution	Scars	
Prevention of the globe				
Multiple operations for cosmetic rehabilitation		Slight scars	Multiple operations for limited visual rehabilitation	




Fig. 2. Grade II chemical injury: lime (CaOH). Central epithelial defect, partial limbal ischemia in the nasal inferior quadrant.

cuers who remove them from dangerous areas and apply fast and efficient help to their eyes and body (Morgan 1987).

Effective first aid involves knowing how to overcome blepharospasm by a passive opening of the lids and how to perform effective irrigation of the eye. All aspects of the conjunctiva and cornea should be irri-

gated, and the patient should be asked to look in all directions (Tannen & Marsden 1991). Topical anesthetic drops may be applied to reduce the pain and to facilitate irrigation. According to the American National Standards Institute (ANSI) standard (Z358.1-1990) severe eye burns have to be rinsed for 15 min. At least 500-1000



Fig. 3. Grade III chemical injury: sodium hydroxide (NaOH). Complete corneal and proximal conjunctival epithelial defect with loss of corneal stromal clarity. Limbal ischemia in the inferior quadrants.

mL of irrigation fluid are thus necessary. Amphoteric or buffered solutions can normalize the pH of the anterior chamber within that time (Schrage et al. 1996). Particles are sometimes trapped in the fornices or under the upper lid. Therefore, ectropinisation and intensive cleaning of the cul-de-sac are mandatory after every burn. Materials containing calcium oxid (lime, cement dust) react avidly with water to produce a calcium hydroxide solution with a pH of 12.4 (Moon & Robertson 1983). A cotton-tipped applicator soaked in EDTA 1% (EDTA, di-sodium-ethylenediaminetetra-acetat) can be used to facilitate cleaning of the cul-de-sac from calcium hydroxide (Pfister 1983). Immediate irrigation is also important in thermal burns, because this cools the ocular surface (Schrage et al. 1997). Continuous irrigation also removes inflammatory substances from the ocular surface (Reim 1990; Reim & Kuckelkorn 1995).

First aid with intensive irrigation immediately after the injury has a decisive influence on the clinical course and prognosis of such eyes (Saari & Parvi 1984; Burns & Paterson 1989). A comparison between visual outcome of better than 1/50 with that of less than 1/50 revealed a highly significant difference, with significantly better results after immediate irrigation. Visual acuity of >1/50 enables the patient to move unaided. The number of operations and the length of stay on the ward are significantly reduced for eyes that received prompt irrigation (Table 3; Kuckelkorn et al. 1995).

Choice of irrigation fluid

Burns are accompanied by a loss of the corneal epithelium within a few seconds. The acutely burnt cornea takes up the burning substance by osmolar forces resulting in a high osmolarity. One of the aims of rinsing therapy is to remove this chemical burden.

Water is commonly recommended as an irrigation fluid. It is available almost everywhere, and copious amounts of water have a dilutive effect. However, water is hypotonic to the corneal stroma and intraocular milieu. In measurements of osmolarity, Schrage et al. (unpublished) found the corneal stroma to have an osmolarity of 420 mOsm/L. The corneal tissue is diluted by rinsing with water and this is accompanied by an increased uptake of additional water and diffusion of the corrosive into the deeper layers of the cornea. We thus recommend the use of irrigation fluids with higher osmolar-

Table 3. Value of immediate rinsing therapy *versus* clinical results in severe eye burns (101 patients, 131 eyes).

Rinsing	Number of operations	Hospitalization (months)	Visual acuity
Immediate	6.5 ± 4.6	4.2 ± 2.8	45 (76%)
Delayed or no	10.4 ± 10	6.0 ± 4.5	22 (55%)

Unpaired *t*-test, *P* < 0.05.

ties for initial rinsing in order to prevent water influx into the cornea and to enable the mobilisation of water and the dissolved corrosives out of the burnt tissue.

Normal saline (NS), which is often recommended as irrigation fluid, also has a lower osmolarity than tear fluid. It fails to normalize the pH of the anterior chamber even after prolonged irrigation (Table 4).

Phosphate buffer is often cited as an

ideal buffer solution (Thiel 1965; Laux et al. 1975; Poser 1983; Roth 1993). For this reason, there is widespread use of this buffer in many factories. However, in an experimental study complete calcification of the superficial stroma occurred in 100% of all animals after burning with 1 n NaOH for 30 s and immediate rinsing with 500 mL phosphate buffer. (Schrage et al., unpublished). We suggest that exogenously applied phosphate reacts with

endogenous calcium released from ruptured cells to produce calcium-phosphate complexes.

At present, there is ongoing experimental research to find irrigation fluids with an osmolarity similar to the corneal stroma. Currently available fluids which are suitable for irrigation are sterile, lactated Ringer's (LR) and balanced saline solution (BSS; Herr et al. 1991). Lactated Ringer is a buffered solution and may be more effective than normal saline. The osmolarity of BSS is similar to that of aqueous humour; its pH is neutral and it contains sodium acetat and citrat (McDermott et al. 1988). According to Pfister, isotonic citrate buffer initiates chelat-complexes and binds unspecific metal-ions derived from the corrosive (Pfister et al. 1981; Pfister et al. 1984). Balanced saline solution has an enhanced buffering capacity; it prevents the cornea from swelling and preserves the corneal endothelium (McNamara et al. 1987).

The pH, osmolarity and buffer capacity of the aqueous humour, corneal stroma and some irrigations fluids are listed in Table 5.

A new amphoteretic solution which is suitable for irrigation is Diphoterine (Previn[®], Fa. Prevor). This newly synthesized fluid is able to bind both alkalis and acids. 0.4% Diphoterine has a pH of 7.4 and an osmolarity of 820 mosm/L. The pH in the conjunctival sac and in the corneal stroma is reduced as rapidly as after irrigation with phosphate buffer. The constituents of Diphoterine are listed in Table 6.

Transport problems

As strong acids and alkalis penetrate within seconds or minutes and remain for hours (Grant & Kern 1955), irrigation should not be interrupted during transport to a professional eye-care unit. The recommendations for minimum irrigation times range from 15 min (ANSI standard; Lubeck & Greene 1988) to 2-4




Fig. 4. Grade IV chemical injury: sodium hydroxide (NaOH). Loss of corneal transparency, ectropion uveae and cataract formation, circular loss of conjunctival and episcleral tissue down to the fornices. The sclera is ischemic.

Table 4. pH on the corneal surface and in the anterior chamber after rinsing with different irrigation fluids (experimental eye burn for 30 s/1 n NaOH).

	pH corneal surface	anterior chamber
Directly after burn	13 ± 0	10 ± 0
5 minutes after rinsing with 500 mL normal saline	9 ± 0	10 ± 0
5 minutes after rinsing with 500 mL phosphat buffer	7.5 ± 0	9.25 ± 0.44
5 minutes after rinsing with 500 mL diphoterine	7.5 ± 0	9.34 ± 0.59

Concentration of phosphat buffer: 440 mg sodiumdihydrogenphosphat and 4040 mg sodiummonohydrogenphosphat in 100 mL H₂O.

Table 5. pH, osmolarity, constituents and buffer capacity of the aqueous humour, corneal stroma and different irrigation fluids.

	pH	osmolarity	constituents	buffer capacity
Aqueous humour	7.4	304	Na, K, Cl, Ca, PO ₄ , proteins	0.0008
Stroma	7.4	420	Na, K, Cl, Ca, S, SO ₄ , PO ₄ , proteins, lipids, glycosaminoglycans	0.0004
Normal saline	7.0	290	Na, Cl	0.0002
Phosphate buffer	7.4	260	Na, K, PO ₄	0.00625
Lactated Ringer's	5.0-7.5	280-309	Na, K, Ca, Cl, lactat	0.00069
BSS	7.2	310	Na, K, Ca, Cl, citrat, acetat	0.001
Diphoterinc	7.4	820	Diphoterinc, Na, Cl, glycin	0.02

h (Pfister 1983; Saari et al. 1984). We recommend the use of an intravenous infusion set to supply at least 500-1000 mL of irrigation fluid. As described above, one of the rescuers should hold the eyelids open while a second rescuer flushes the eye with a mild, directable and controllable stream of fluid. Local anesthetic drops should be administered repeatedly if necessary to relieve the patient from pain and to facilitate irrigation.

Some authors favor the use of specially designed irrigation systems (Naumann 1964; Girard & Soper 1966; Schulze & Tost 1967; Tan 1970; Morgan 1971; Lau 1979). Whereas these systems provide continuous irrigation of the eye, they fail to flush the ocular surface homogenously and appropriately, especially the cul-de-sac. A further risk with lime or cement burnsis that particles retained under the eye lids are not detected and removed once the loop or lens has been applied. Moreover, rescuers unfamiliar with the handling of these systems will lose precious time when installing the slings or lenses, which may cause additional damage to the eye.

The effectiveness of rinsing therapy can be assessed by using universal indicator paper to determine the pH of the external eye. Irrigation must be continued as long as the pH remains outside the normal range. If prolonged irrigation does not achieve normalization of the pH, one

must consider the possibility that there are still particles in the superior or inferior cul-de-sac.

Subsequent care

The subsequent care of eye burns is dependent on the severity of the injury. Further therapeutical procedures are applied according to the extent of the damage. If the injury is mild (grades I and II) and irrigation began immediately, most eyes will heal without permanent damage within a few days (Moon & Robertson 1983; Morgan 1987; Beare 1990; Kuckelkorn et al. 1993). Topical steroid/antibiotic drops and ointment plus padding may suffice for the treatment of these mild burns. Follow-up treatment within 24h is mandatory.

Severe ocular eye burns (grades III and IV) are difficult to treat and the course of healing often takes several months. In these cases, accurate classification with regard to the extend of (limbal) ischemia and depth of tissue destruction is essential. An examination with the operating microscope is thus mandatory. Parabulbar or general anesthesia are sometimes needed if the patient suffers pain and local anesthetic drops are not sufficient.

Less severe eye burns (grade III) are characterized by superficial ischemia of the conjunctival tissue. In these cases where regular anterior chamber structures are preserved and there is no damage to the iris, ectropium uveae or fibrinous exsudation, subsequent management takes the form of a more conservative therapy. Admission to and treatment in a local eye clinic are thus sufficient (Reim & Kuckelkorn 1995).

Most severe eye burns (grade IV) lead to significant limbal ischemia and necrosis of the bulbar and tarsal conjunctiva as well as of the episcleral tissue

down to the fornices. In cases with superficial necrosis, the deep episcleral vessels are still perfused while necrosis of the deeper episcleral tissue is associated with thrombosis of the episcleral vessels. In these severe cases, opacification of the cornea is common and the anterior chamber structures are obscured. A greyish aspect of the iris, ectropium uveae and the breakdown of the blood aqueous barrier with fibrinous exsudation into the anterior chamber confirm the destruction of the deep anterior segment. The lids and aspects of the tarsal conjunctiva are often involved. Many problems arise in the acute phase of the burn. The most delicate problem is preventing the eye from early melting.

Necrosis of the conjunctiva and subconjunctival tissue is accompanied by a considerable exsudation of leucocytes (PMN's). These leucocytes release large amounts of lysosomal enzymes. The matrix metalloproteinases (MMP); collagenase (MMP-1 and MMP-8; Itoi et al. 1969; Hook et al. 1971; Newsome & Gross 1977; Johnson-Muller & Gross 1978; Kuter et al. 1989; Fini & Girard 1990), gelatinase (MMP-2, MMP-9) (Collier et al. 1988; Fini & Girard 1990; Huhtala et al. 1990), and stromelysin (MMP-3; Collier et al. 1988; Chin et al. 1995) in particular are responsible for the splitting of the collagen molecules and the development of corneoscleral and corneal ulceration, characteristically 4-6 weeks after the accident.

The basic principle in the treatment of these eyes is to reduce the inflammatory response caused by the necrotic tissue. The traditional mainstay of therapy is the early and intensive application of corticosteroids (Donshik et al. 1978; Leibowitz 1980; Reim & Schmidt-Martens 1982; Kenyon 1985; Reim 1987). Additionally, local antibiotics are necessary to prevent microbiobal infections until the ocular surface has reepithelialized (Girard et al. 1970; Kuckelkorn et al. 1987; Beare 1990). Tetracycline derivates play an important role because they have been shown to inhibit metallo-proteinases (Brion et al. 1985; Golup et al. 1987; Seedor et al. 1987; Burns et al. 1989; Perry et al. 1993) independently of their anti-microbial properties.

Besides conservative therapy, active surgical intervention with the débridement of necrotic conjunctival and subconjunctival tissue is necessary in order to remove a nidus of continued inflammation from retained caustic materials,

Table 6.

Constituents of diphoterinc.	
Amphoter: Diphoterinc	3.8/100mL
NaCl:	1.8/100mL
Glycin:	0.75/100mL
Preservative:	0.05/100mL
Aqua destillata:	ad 100mL

although also any accumulation of PMN and to prevent the sustained release of their destructive enzymes. Special reconstructive approaches such as tenonplasty allow the denuded avascular sclera to be covered with vital connective tissue prepared from the equator of the globe (Reim & Teping 1989; Reim 1992; Reim & Kuckelkorn 1992, 1995; Kuckelkorn & Reim 1993; Reim & Leber 1993; Kuckelkorn et al. 1995). The main advantage of this tissue is that it enables the reconstruction of the conjunctival matrix and of limbal vascularity. These interventions prevent anterior segment necrosis and/or sterile ulceration and the eye is preserved. It is advisable to admit these cases to an eye clinic specialized in the treatment of these eyes and familiar with specialized procedures in plastic reconstruction.

References

- Ballen PH (1963): Mucous membrane grafts in chemical (lye) burns. *Am J Ophthalmol* **55**: 302-312.
- Beare JDL (1990): Eye injuries from assault with chemicals. *Br J Ophthalmol* **74**: 514-518.
- Brion M, Lambs L & Berthon G (1985): Metal ion-tetracycline interactions in biological fluids: Formation of zinc complexes with tetracycline and some of its derivatives and assessment of their biological significance. *Agents Actions* **17**: 229-242.
- Burns FR & Paterson CA (1989): Prompt irrigation of chemical eye injuries may avert severe damage. *Occup Health Saf* **58**: 33-36.
- Burns FR, Stack M & Gray R (1989): Inhibition of purified collagenase from alkali-burned rabbit corneas. *Invest Ophthalmol Vis Sci* **30**: 1569-1575.
- Chin J, Murphy G & Werb Z (1995): Stromelysin, a connective-tissue degrading metalloendopeptidase secreted by rabbit synovial fibroblasts in parallel with collagenase. *J Biol Chem* **260**: 12367-12376.
- Cohen KL & Hyndiuk RA (1978): Ocular emergencies. *Am Fam Physician* **18**: 178-184.
- Collier I, Smith J & Kronberger A (1988): The structure of the human skin fibroblast collagenase gene. *J Biol Chem* **263**: 10711-10713.
- Collier I, Wilhelm S & Eisen A (1988): II-ras oncogene-transformed human bronchial epithelial cells (TBE-1) secrete a single metalloprotease capable of degrading basement membrane collagen. *J Biol Chem* **263**: 10711-10713.
- Doushik PC, Berman MB, Dohlman CH et al. (1978): Effect of topical corticosteroids on ulceration in alkali-burned corneas. *Arch Ophthalmol* **96**: 2117-2120.
- Eakins K & Bhattacharjee P (1977): Histamine, prostaglandins and ocular inflammation. *Exp Eye Res* **24**: 299-305.
- Fini M & Girard M (1990): Expression of collagenolytic/gelatinolytic metalloproteinases by normal cornea. *Invest Ophthalmol Vis Sci* **31**: 1779-1788.
- Friedenwald J, Hughes W & Hermann H (1944): Acid-base tolerance of the cornea. *Arch Ophthalmol* **31**: 279-283.
- Friedenwald JS, Hughes WF & Hermann H (1946): Acid burns of the eye. *Arch Ophthalmol* **35**: 98-108.
- Girard LJ, Alford WE, Feldman GL & Williams B (1970): Severe alkali burns. *Trans Am Acad Ophthalmol Otol.*
- Girard LJ & Soper J (1966): Flush-fitting scleral contact-lenses. *Am J Ophthalmol* **61**: 169-183.
- Golup L, McNamara T, D'Angelo G & Greenwald RA (1987): A non antibacterial chemically-modified tetracycline inhibits mammalian collagenase activity. *J Dent Res* **66**: 1310-1314.
- Grant WM (1974): Toxicology of the Eye. In: Charles C Thomas (ed.) *Drugs, Chemicals, Plants, Venoms*. Springfield: 5-22 and 96-101.
- Grant W & Kern H (1955): Action of alkalis on the corneal stroma. *Arch Ophthalmol* **54**: 931-939.
- Graupner O & Hausmann H (1970): Die Änderung des pH-Wertes in der Vorderkammer des Kaninchenauges nach Verätzung mit kleinsten Mengen laborüblicher Konzentrationen von Säure und Lauge. *Albrecht Von Graefes Arch Clin Exp Ophthalmol* **180**: 60.
- Herr R, White G, Bernhisel K, Mamalis N & Swanson E (1991): Clinical comparison of ocular irrigation fluids following chemical injury. *Am J Emerg Med* **9**: 228-231.
- Hook C, Brown S, Iwanji W & Nakanishi I (1971): Characterization and inhibition of corneal collagenase. *Invest Ophthalmol Vis Sci* **10**: 496-503.
- Hughes WJ (1946): Alkali burns of the eye. II. Clinical and pathological course. *Arch Ophthalmol* **36**: 189-214.
- Huhtala P, Chow I & Tryggvason K (1990): Structure of the human type IV collagenase gene. *J Biol Chem* **265**: 11077-11082.
- Itoi M, Gnädiger M & Slansky H (1969): Collagenase in the cornea. *Exp Eye Res* **8**: 369.
- Johnson-Muller B & Gross J (1978): Regulation of corneal collagenase production: epithelial stromal cell interactions. *Proc Natl Acad Sci USA* **75**: 4417-4421.
- Keeney A (1974): Prevention of ocular injuries. *Int Ophthalmol Clin* **14**: 1-10.
- Kenyon KR (1985): Inflammatory mechanisms in corneal ulceration. *Trans Am Ophthalmol Soc* **83**: 610-663.
- Kuckelkorn R (1996): Epidemiologie und chirurgische Behandlung schwerster Verätzungen und Verbrennungen der Augen. *Habilitationsschrift*. Thesis.
- Kuckelkorn R, Kottek A, Schrage N & Reim M (1995): Poor prognosis of severe chemical and thermal burns. The need for adequate emergency care and primary prevention. *Int Arch Occup Environ Health* **67**: 281-284.
- Kuckelkorn R, Kuwert T, Melzer T & Reim M (1987): Hornhautbefunde bei schwersten Verätzungen und Verbrennungen der augen. *Fortschr Ophthalmol* **84**: 540-544.
- Kuckelkorn R, Luft I, Kottek A, Schrage N, Makropoulos W & Reim M (1993): Verätzungen und Verbrennungen im Einzugsbereich der RWTH Aachen. *Klin Monatsbl Augenheilkd* **203**: 34-41.
- Kuckelkorn R, Makropoulos W, Kottek A & Reim M (1995): Chemische Verätzungen der augen: ursachen, behandlung und prävention. *Arbeitsmed Sozialmed Umweltmed* **30**: 22-33.
- Kuckelkorn R, Redbrake C, Kottek A, Feldmann P & Reim M (1995): Tenon-plastik und frühkeratoplastik bei schwerstverätzten augen. *Ophthalmologie* **92**: 439-444.
- Kuckelkorn R & Reim M (1993): Wiederherstellungsoperation nach schwerster Verätzung mit perforation. *Fortschr Ophthalmol* **90**: 716-719.
- Kuckelkorn R, Schrage N & Reim M (1995): Treatment of severe eye burns by tenonplasty. *Lancet* **345**: 657-658.
- Kulkarni P & Srinivasan B (1983): Synthesis of slow reacting substance-like activity in rabbit conjunctiva and anterior uvea. *Invest Ophthalmol Vis Sci* **24**: 1079-1085.
- Kuter I, Johnson-Wint B, Beupre N & Gross J (1989): Collagenase secretion accompanying changes in cell shape occurs only in the presence of a biologically active cytokine. *J Cell Sci* **92**: 473-492.
- Lau L (1979): Continuous lavage in (chemical) corneal lesions. *Doc Ophthalmol* **46**: 215-218.
- Laux U, Roth H & Steinhardt B (1975): Die Wasserstoffionen-konzentration des Kammerwassers nach Alkaliverletzung der Hornhaut und deren therapeutische Beeinflussbarkeit. *Graefes Arch Clin Exp Ophthalmol* **195**: 33-40.
- Leibowitz HM (1980): Management of inflammation in the cornea and conjunctiva. *Ophthalmology* **87**: 753-758.
- Liggett P (1989): Ocular trauma in an urban population. *Ophthalmology* **97**: 581-584.
- Lubeck D & Greene J (1988): Corneal injuries. *Emerg Med Clin North Am* **6**: 73-94.
- MacEwen C (1989): Eye injuries: a prospective survey of 5,671 cases. *Br J Ophthalmol* **73**: 888-894.
- McCulley JP (1987): Chemical injuries. In: Smolin, G, Thoft, RA (eds.): *The Cornea*. Boston. Little Brown: 527.
- McDermott MI, Edelhauser HF, Hack HM et al. (1998): Ophthalmic irrigants. A current review and update. *Ophthalmic Surg* **19**: 724-733.
- McNamara MJ, Seaber AV & Urbaniak JR (1987): Effect of irrigation fluids on arterial

- and venous endothelium after ischemia. *J Reconstr Microsurg* **4**: 27-32.
- Moon MEL & Robertson IF (1983): Retrospective study of alkali burns of the eye. *Aust J Ophthalmol* **11**: 281-286.
- Morgan I. (1971): A new drug delivery system for the eye. *Industrial Med* **40**: 11-13.
- Morgan S (1987): Chemical burns of the eye: causes and management. *Br J Ophthalmol* **71**: 854-857.
- Morris R, Witherspoon C & Holms HJ (1987): Eye injury Registry of Alabama (preliminary report): demographics and prognosis of severe eye injury. *South Med J* **80**: 810-881.
- Naumann (1964): Ein einfacher vorschlag zur behandlung des vorderen augenabschnittes im dauerdropfverfahren. *Klin Monatsbl Augenheilkd* **144**: 597-600.
- Newsome N & Gross J (1977): Prevention by medroxyprogesteron of perforation in the alkali-burnt rabbit cornea: inhibition of collagenolytic activity. *Invest Ophthalmol Vis Sci* **16**: 21-31.
- Perry H, Hodes L & Seedor J (1993): Effect of doxycycline hyclate on corneal epithelial wound healing in the rabbit alkali-burn model. Preliminary observations. *Cornea* **12**: 379-382.
- Pfister R (1983): Chemical injuries of the eye. *Ophthalmology* **90**: 1246-1253.
- Pfister R, Haddox J, Dodson RW & de Shazo W (1984): Polymorphonuclear leukocytic inhibition by citrate, other metal chelators and trifluoprazine: evidence to support calcium binding protein involvement. *Invest Ophthalmol Vis Sci* **25**: 79.
- Pfister R, Nicolario M & Paterson CA (1981): Sodium citrate reduces the incidence of corneal ulcerations and perforations in extreme alkali-burned eyes-acetylcysteine and ascorbat have no favorable effect. *Invest Ophthalmol Vis Sci* **21**: 486-490.
- Poser E (1983): The emergency therapy of burns. *IMJ* **164**: 186-187.
- Reim M (1982): The clinical significance of proteolytic enzymes and glycosidases in corneal disease. *Trans Asian Pacific Acad Ophthalmol* **3**: 342-346.
- Reim M (1987): Zur behandlung von schwersten verätzungen und verbrennungen der bindehaut. *Fortschr Ophthalmol* **84**: 46-54.
- Reim M (1987): Zur pathophysiologie und therapie von verätzungen. *Fortschr Ophthalmol* **84**: 65-69.
- Reim M (1990): Ein neues behandlungskonzept für schwere verätzungen und verbrennungen der augen. *Klin Monatsbl Augenheilkd* **196**: 1-5.
- Reim M (1992): The results of ischemia in chemical injuries. *Eye* **6**: 376-380.
- Reim M, Bahrke C, Kuwert T & Kuckelkorn R (1993): Investigation of enzyme activities in severe burns of the anterior eye segment. *Gräfe's Arch Clin Exp Ophthalmol* **231**: 308-312.
- Reim M & Kuckelkorn R (1992): Chemical and thermal lesions in the orbital region. *Bull Soc Belge Ophthalmol* **245**: 21-28.
- Reim M & Kuckelkorn R (1995): Verätzungen und verbrennungen der augen. *Akt Augenheilkd* **20**: 76-89.
- Reim M & Leber M (1993): N-Acetylglucosaminidase activity in corneoscleral ulceration after severe eye burns. *Cornea* **12**: 1-7.
- Reim M & Schmidt-Martens FW (1982): Behandlung von verätzungen. *Klin Monatsbl Augenheilkd* **181**: 1-9.
- Reim M & Teping C (1989): Surgical procedures in the treatment of severe eye burns. *Acta Ophthalmol Suppl* **67**: 47-54.
- Rochels R, Busse W, Haekelbusch R & Knieper P (1982): Prostaglandin-konzentrationsbestimmung im kammerwasser nach der bindehaut-hornhautverätzung: korrelation biochemischer, funktioneller und morphologischer befunde. *Fortschr Ophthalmol* **80**: 145-147.
- Rodeheaver GT, Hiebert JM & Edlich RF (1982): Initial treatment of chemical skin and eye burns. *Compre Ther* **8**: 37-43.
- Roper-Hall MJ (1965): Thermal and chemical burns. *Trans Ophthalmol Soc UK* **85**: 631-653.
- Roth H (1993): Untersuchungen zur sofortbehandlung von augenverätzungen mit einem phosphatpuffer. *Augenspiegel* **39**: 6-11.
- Saari K, Leinonen J & Aine E (1984): Management of chemical eye injuries with prolonged irrigation. *Acta Ophthalmol Suppl* **161**: 52-59.
- Saari K & Parvi V (1984): Occupational eye injuries in Finland. *Acta Ophthalmol Suppl* **161**: 17-28.
- Schrage N, Flick S & Aschenbrenner W (1996): Rinsing therapy in severe alkali burns of rabbit eyes. *Vision Res Supplement* **1**: 248.
- Schrage NF, Flick S & Fischern von T (1997): How do different types of bandages influence the corneal temperature. *Ophthalmologie* **94**: 492-495.
- Schulze F & Tost M (1967): Neue applikatoren zur dauerdropftherapie in der ophthalmologie und ihre anwendungsmöglichkeiten. *Klin Monatsbl Augenheilkd* **151**: 471-477.
- Seedor J, Perry H & McNamara T (1987): Systemic tetracycline treatment of alkali-induced corneal ulceration in rabbits. *Arch Ophthalmol* **105**: 268-271.
- Struck H, Geiser H & Taube C (1991): The antiphlogistic effect of the ginkgolide BN 52021 in the chemically burned rabbit eye. *Klin Monatsbl Augenheilkd* **199**: 278-282.
- Tan B (1970): Oklahoma eye irrigation tube. *Trans Am Acad Ophthalmol Otolaryngol* **74**: 435-437.
- Tannen M & Marsden J (1991): Chemical burns of the eye. *Nursing Standard* **6**: 24-26.
- Thiel R (1965): Behandlung von verätzungen. *Klin Monatsbl Augenheilkd* **146**: 581-587.
- Thielsch J, Parver L & Shankar B (1989): Time trends in the incidence of acute hospitalized ocular trauma. *Arch Ophthalmol* **107**: 519-523.
- Watz H & Reim M (1973): Aus der unfallstatistik einer ländlichen augenklinik. *Klin Monatsbl Augenheilkd* **162**: 648-655.
- Williams R, Bhattacharjee P & Eakins K (1983): Biosynthesis of lipoxigenase products by ocular tissues. *Exp Eye Res* **36**: 397-402.
- Zagelbaum B, Tostanowski J, Kerner D & Hersh P (1993): Urban eye trauma. *Ophthalmology* **100**: 851-856.

Received on April 9th, 2001.

Accepted on September 9th, 2001.

Correspondence:

Ralf Kuckelkorn, MD

Wilhelmstrasse 8

52070 Aachen

Germany